

STATE OF THE ART AND SCIENCE

Continuing Medical Education and Firearm Violence Counseling

Nicole D. Damari, Karan S. Ahluwalia, Anthony J. Viera, MD, MPH, and Adam O. Goldstein, MD, MPH

Abstract

Firearm violence is a significant and increasing cause of mortality. Although physicians view firearm counseling as their professional obligation, few engage in the practice. This study examines medical education and firearm counseling among physicians in North Carolina. While 65 percent of physicians reported knowing how to counsel patients about gun safety, only 25 percent reported having conversations with patients about firearms or firearm safety often or very often. Physicians reporting continuing medical education (CME) attendance on gun safety, however, were more likely to report providing patients with firearm safety counseling and asking patients with depression about firearms. Increasing availability of and physician participation in firearm violence prevention CME could significantly increase physicians' knowledge of and engagement in firearm counseling.

Introduction

Firearm violence is a significant cause of mortality that has increased as an important ethical and public health issue. Data from 2014 revealed over 33,000 firearm-related deaths in the United States, accounting for 16.8 percent of all injury-related deaths, 49.6 percent of all completed suicides, and 69.2 percent of homicides [1]. In 2014, North Carolina, the state in which the authors reside, ranked seventh among states [2], with 1,146 firearm deaths, representing 59.3 percent of all violent death in the state that year [3].

Many medical organizations and physicians have discussed the [roles clinicians should play](#) in preventing firearm injury and deaths [4-13]. Surveys of family physicians, internists, psychiatrists, pediatricians, and surgeons have revealed that between 65 to 93 percent believe gun safety counseling is within a physician's scope of practice [14-20]. Many physicians also believe firearm safety counseling is effective at reducing rates of firearm-related suicides and homicides [21, 22].

Although physicians view such counseling as part of their professional obligations, few appear to engage in the practice. Chart reviews from internal medicine and pediatric

emergency departments demonstrate that 3 percent or less document gun access or [firearm safety counseling](#) [23, 24]. While chart review may not capture every service delivered, documentation likely serves as proxy for what physicians believe is important from treatment planning and medical-legal perspectives [24]. A survey of 573 internists revealed that 58 percent had never asked patients about gun ownership, and 77 percent had never counseled patients on risks of firearm-related injury or death [20]. Similarly, less than half of surveyed pediatricians reported regularly providing firearm safety screening or counseling [18, 25-27]. At the medical student level, 66 percent of 1,469 seniors across 16 schools reported counseling patients about firearm safety “never” or “rarely,” with only 4 percent reporting doing so “usually” or “always” [16].

Discrepancies between medical professional beliefs and behaviors with respect to firearm safety counseling signal an opportunity for intervention should their cause be identified. Existing literature suggests that physician training in firearm safety education is woefully inadequate [16, 18, 20-22, 28]. When asked directly whether there was a need for physician education on firearm injury prevention counseling, 74 percent of 573 internists answered “somewhat” or “to a great extent” [20]. While two-thirds of 1658 medical students reported receiving any training on firearm safety counseling, only 12 percent considered the training “extensive” [16]. At the residency level, a survey of pediatric residency programs revealed that only one-third include formal training on firearm safety counseling [18]. Similarly, among psychiatric residency directors, 79 percent of respondents had not considered including training to address firearm injury prevention, citing lack of standardized material, faculty expertise, and training guidelines as key barriers [21], echoing the sentiments of preventive medicine programs [22]. A recent review found that only four firearm safety training programs report targeting medical education, none of which targeted psychiatrists or psychiatry residents [28].

While the existing literature addresses the need for firearm safety training in undergraduate and graduate medical education, little research has been conducted on the role of continuing medical education (CME) for physicians about firearm violence prevention counseling. This survey-based study of North Carolina internists, psychiatrists, and family medicine physicians examines how various criteria, including CME, affect firearm safety counseling confidence and behaviors for clinicians in practice. Other questions that examined physician experiences with and attitudes toward concealed weapon permits, which showed that physicians were often unsure of and had little confidence in determining competence for concealed weapons’ permitting, were previously published [29, 30].

Methods

Survey. The survey questions addressed physician attitudes, beliefs, and behaviors about various issues related to firearm safety, including patient counseling and self-reported CME on firearm safety education. Physician attitudes and beliefs were assessed by four

categories of response (strongly agree, agree, disagree, and strongly disagree), and physician behaviors were assessed by five categories of response (never, rarely, sometimes, often, and very often). Respondent demographic information included age, sex, specialty, political affiliation, years in practice, gun ownership, and experience shooting a gun.

Participants. The survey was mailed to 600 physicians registered with the North Carolina Medical Board and in active practice at the time of the survey (September and October of 2013), including 200 physicians identified as family physicians, psychiatrists, and internists who were randomly selected to receive the survey. Participants in the second and third of three survey waves received nominal incentives (<\$1) to participate. Of the 600 surveys mailed, 45 were returned because of incorrect addresses and 223 were completed, for an adjusted response rate of 40.2 percent. The pool of respondents included psychiatrists (38 percent), family physicians (35 percent), and internists (27 percent). The majority of respondents were men (66 percent), over age 51 (55 percent), and had been in practice more than 15 years (64 percent). Most self-described as Democrat (47 percent), with the remaining being Republican (20 percent), Independent (25 percent), or "other," including libertarian (8 percent). Seventy-six percent of physicians in the sample reported having shot a gun, and 36 percent identified as gun owners.

Analyses. The Pearson chi-square test was used to examine the relationships between survey response variables and responses to three of the attitude and behavior questions, with significance judged by Bonferroni corrected alpha levels of 0.006 (0.05/9) or 0.005 (0.05/10). For attitude and belief questions, agree/strongly agree and disagree/strongly disagree were combined to yield two response categories. Similarly, for behavior questions, never/rarely and often/very often were combined to yield three responses categories. Logistic regression was used to estimate the odds of associations between CME and aspects of gun safety counseling.

Results

Descriptive statistics. Most physicians agreed or strongly agreed that gun violence was a major public health issue that should be part of medical training (80 percent). Almost two-thirds (65 percent) of physicians reported knowing how to counsel patients about gun safety, but only 25 percent reported having conversations with patients about firearms or firearm safety often or very often. With regard to suicide prevention, only 52 percent of physicians reported asking depressed patients if they had a firearm in their home. Furthermore, only 12 percent of physicians reported having attended any continuing medical education (CME) seminars or lectures on gun violence in the last five years. CME attendance was not related to political party affiliation.

Factors associated with knowledge of gun safety counseling. As shown in Table 1, physicians' self-reported knowledge of how to counsel patients about gun safety was significantly related to physician specialty, with psychiatrists most likely to report knowledge of how to counsel about gun safety compared to family physicians and internists (78.6 percent vs. 66.7 percent vs. 45.0 percent respectively; $p < 0.001$). Those who reported attending CME on gun safety were more likely to say they knew how to counsel about gun safety, but this was of borderline significance (88.9 percent vs. 61.7 percent; $p = 0.006$). Sex, age, years in practice, political affiliation, gun ownership, and having shot a gun were not significantly associated with knowledge of gun safety counseling.

Table 1. Bivariate associations for “knowing how to counsel patients about gun safety”

Variable	Agree/strongly agree (%)	<i>p</i> -value ^a
Sex (<i>N</i> = 222)		
Male	68.5	0.17
Female	59.2	
Specialty (<i>N</i> = 222)		
Family medicine	66.7	< 0.001
Psychiatry	78.6	
Internal medicine	45.0	
Age (<i>N</i> = 222)		
< 35	51.2	0.09
36-50	74.0	
> 51	63.1	
Years in practice (<i>N</i> = 222)		
< 5	55.2	0.46
5-15	68.2	
> 15	66.4	
Political affiliation (<i>N</i> = 220)		
Democrat	56.7	0.04
Republican	70.5	
Independent	78.3	
Other	70.6	
Shot a gun (<i>N</i> = 222)		
Yes	65.7	0.84
No	64.2	

Owns a gun (<i>N</i> = 222)		
Yes	75.0	
No	59.9	0.02
Gun violence should be part of medical student training (<i>N</i> = 219)		
Agree/strongly agree	67.4	
Disagree/strongly disagree	61.4	0.45
CME attendance (<i>N</i> = 223)		
Yes	88.9	
No	61.7	0.006

^a Bonferroni corrected alpha level = 0.006.

Factors associated with counseling frequency. As shown in Table 2, reported knowledge of gun safety counseling was significantly related to reported frequency of counseling about gun safety (33.1 percent vs. 10.3 percent counseling very often/often; $p < 0.001$). Frequency of counseling differed significantly by specialty, with psychiatrists reporting higher rates of counseling “very often or often” compared to family physicians and internists (48.8 percent vs. 15.0 percent vs. 7.7 percent; $p < 0.001$). Those who reported attending CME on gun safety counseling were more likely to say they counseled patients often or very often compared to those who did not attend CME (59.3 percent vs. 20.4 percent; $p < 0.001$). Reports on counseling frequency were not associated with sex, age, years in practice, political affiliation, gun ownership, or having shot a gun.

Table 2. Bivariate associations for “counseling frequency about gun safety”

Variable	Never/ rarely (%)	Some- times (%)	Often/very often (%)	<i>p</i> -value ^a
Sex (<i>N</i> = 222)				
Male	50.7	27.4	21.9	
Female	36.8	31.6	31.6	0.12
Specialty (<i>N</i> = 222)				
Family medicine	59.0	33.3	7.7	
Psychiatry	25.0	26.2	48.8	
Internal medicine	58.3	26.7	15.0	< 0.001
Age (<i>N</i> = 222)				
< 35	62.9	18.5	18.5	
36-50	32.9	37.0	30.1	
> 51	50.0	26.2	23.8	0.06

Years in practice (<i>N</i> = 222)				
< 5	55.2	20.7	24.1	
5-15	32.0	38.0	30.0	
> 15	49.0	27.3	23.8	0.22
Political affiliation (<i>N</i> = 220)				
Democrat	46.2	28.9	25.0	
Republican	54.6	22.7	22.7	
Independent	38.2	32.7	29.1	
Other	41.2	35.3	23.5	0.80
Shot a gun (<i>N</i> = 222)				
Yes	46.8	30.8	22.5	
No	43.4	22.6	34.0	0.21
Owns a gun (<i>N</i> = 222)				
Yes	46.3	33.8	20.0	
No	45.8	26.1	28.2	0.30
Gun violence should be part of medical student training (<i>N</i> = 219)				
Agree/strongly agree	41.7	29.7	28.6	
Disagree/strongly disagree	63.6	22.7	13.6	0.03
I know how to counsel patients about gun safety (<i>N</i> = 223)				
Yes	36.6	30.3	33.1	
No	64.1	25.6	10.3	< 0.001
CME attendance (<i>N</i> = 223)				
Yes	14.8	25.9	59.3	
No	50.5	29.1	20.4	< 0.001

^a Bonferroni corrected alpha level = 0.005.

Factors associated with asking depressed patients about firearm access. Factors associated with reports on the frequency of asking depressed patients about access to firearms are shown in Table 3. Asking depressed patients about access to firearms was positively associated with reported knowledge of how to counsel (58.6 percent vs. 39.7 percent; $p < 0.001$), specialty (72.6 percent vs. 41.0 percent vs. 38.3 percent; $p < 0.001$), and CME attendance on gun counseling (85.2 percent vs. 47.5 percent; $p < 0.001$). Reports on counseling frequency were not associated with sex, age, years in practice, political affiliation, gun ownership, or having shot a gun.

Table 3. Bivariate associations for “asking depressed patients about access to firearms”

Variable	Never/ rarely (%)	Some- times (%)	Often/very often (%)	<i>p</i> -value ^a
Sex (<i>N</i> = 222)				
Male	28.8	19.2	52.1	0.62
Female	23.7	23.7	52.6	
Specialty (<i>N</i> = 222)				
Family medicine	32.1	26.9	41.0	< 0.001
Psychiatry	9.5	17.9	72.6	
Internal medicine	45.0	16.7	38.3	
Age (<i>N</i> = 222)				
< 35	25.9	29.6	44.4	0.78
36-50	27.4	17.8	54.8	
> 51	27.1	20.5	52.5	
Years in practice (<i>N</i> = 222)				
< 5	27.6	27.6	44.8	0.25
5-15	20.0	14.0	66.0	
> 15	29.4	21.7	49.0	
Political affiliation (<i>N</i> = 220)				
Democrat	25.0	23.1	51.9	0.54
Republican	36.4	18.2	45.5	
Independent	23.6	21.8	54.6	
Other	17.7	11.8	70.6	
Shot a gun (<i>N</i> = 222)				
Yes	26.6	18.3	55.0	0.22
No	28.3	28.3	43.4	
Owns a gun (<i>N</i> = 222)				
Yes	28.8	22.5	48.8	0.73
No	26.1	19.7	54.2	
Gun violence should be part of medical student training (<i>N</i> = 219)				
Agree/strongly agree	22.3	21.7	56.0	0.007
Disagree/strongly disagree	45.5	18.2	36.4	

I know how to counsel patients about gun safety (<i>N</i> = 223)				
Yes	19.3	22.0	58.6	
No	42.3	18.0	39.7	0.001
CME attendance (<i>N</i> = 223)				
Yes	3.7	11.1	85.2	
No	30.6	21.9	47.5	0.001

^a Bonferroni corrected alpha level = 0.005.

Association between CME attendance and outcomes. After adjustment for specialty, gun ownership, and political party, physicians reporting CME attendance had 3.2 times the odds of reporting providing patients with gun safety counseling (aOR 3.23; 95 percent CI 1.2-8.5; $p = 0.018$) and 4.4 times the odds of reporting asking depressed patients about firearms (aOR 4.37; 95 percent CI 1.38-13.8; $p = 0.012$). CME attendees also had 3 times the odds of reporting knowledge of how to counsel (aOR 3.01; 95 percent CI 0.84-10.8; $p = 0.092$), although this last result had a wide confidence interval and was not statistically significant.

Discussion

It is increasingly clear that addressing firearm safety in clinical situations constitutes an ethical imperative that physicians should satisfy, particularly as physician skills in counseling patients about firearm violence prevention are increasingly considered key to helping prevent firearm deaths [12, 31]. Fulfilling this imperative requires the cultural competence and nuance necessary to have dedicated and respectful conversations with patients [13]. While low rates of gun safety counseling and lack of physician training to provide counseling are established [6, 12, 28, 30, 32], the present research is the first to illuminate the potent role that CME may play in increasing counseling. CME was strongly associated with providing firearm counseling often or very often, including asking patients with depression about their firearm access as well as increased knowledge of how to counsel.

Just as educational experiences have been shown to positively impact physician knowledge of addressing adolescent violence or geriatric suicide risk [15, 33-35], CME likely provides a critical opportunity for practicing physicians to acquire ongoing practical skills that significantly improve their firearm counseling behaviors. While the causal relationship could be reversed, with physicians who already prioritize gun safety seeking related CME, it seems unlikely that physicians who already have significant gun safety knowledge and counseling skills would invest in CME on that topic. Furthermore, since 80 percent of respondents in the current study agreed or strongly agreed that gun violence is a major public health problem that should be part of medical education, a

large gap exists in skill levels between the minority of respondents who do regularly provide gun safety counseling and the great majority of respondents who do not do so.

Our findings extend the literature on physicians' role in addressing firearm violence spanning individual specialties and geographic locations [18, 20, 29, 30, 32], providing a glimpse of the current landscape on primary care and specialty involvement in firearm counseling practices. Psychiatrists in our sample appeared to report higher knowledge of how to counsel, higher frequency of counseling, and asking patients with depression more often about access to firearms. Three-quarters of psychiatrists did endorse that they possessed knowledge of how to counsel patients, indicating that psychiatrist training or practice breeds more knowledge of counseling behaviors.

The ethical imperative to not neglect firearm safety in clinical situations is particularly relevant to suicide prevention, as suicide by firearms accounts for one-half of recorded suicides, and depression is strongly linked with suicidality [2]. Intra-specialty analysis reveals that few family physicians (7.7 percent) or internists (15 percent) reported asking patients with depression about firearm access either very often or often. Even among psychiatrists, who were significantly more likely to report asking their depressed patients about firearm access, less than half reported counseling patients with depression about their firearm access.

Physicians' provision of gun safety counseling is often highlighted as a polarizing political issue, particularly in light of legislative attempts to block physicians from discussing firearm violence prevention with patients [36]. However, our research counters this narrative, as neither political party nor gun ownership was significantly associated with self-reported counseling knowledge, counseling frequency, or rates of asking patients with depression about firearm access. These findings are promising, as they suggest that the provision of firearm safety counseling need not be a partisan issue; instead, our findings provide an opportunity for physicians across party lines to sensibly unite behind the need to provide accurate and effective firearm counseling to patients, as has been done before with safety issues like domestic violence, seatbelts, biking helmets, and smoking cessation. Perhaps the disparity between belief and low CME attendance rate is not driven by a lack of interest or perceived benefit but by the paucity of available CME firearm safety opportunities [28].

Conclusion

While research to replicate and extend our findings to other physician populations and states is needed, our study suggests that increasing availability of, and physician enrollment in, firearm safety-focused CME could impact knowledge of and counseling on firearm violence prevention. Firearm safety educational interventions could empower the great majority of physicians who have not had exposure to this type of training to

provide vital safety counseling for their patients, ultimately contributing to fulfillment of physicians' obligations to help reduce firearm injury and death.

References

1. Kochanek KD, Murphy SL, Xu J, Tejada-Vera B. Deaths: final data for 2014. *Natl Vital Stat Rep*. 2016;65(4):1-122.
2. Centers for Disease Control and Prevention National Center for Health Statistics. Firearm mortality by state: 2014. https://www.cdc.gov/nchs/pressroom/sosmap/firearm_mortality/firearm.htm. Accessed November 20, 2017.
3. North Carolina Department of Health and Human Services Injury and Violence Prevention Branch. Firearm deaths in North Carolina, 2014. http://www.injuryfreenc.ncdhhs.gov/DataSurveillance/VDRS/NC-VDRSFirearmDeaths2014_14Dec16.pdf. Published December 2016. Accessed September 18, 2017.
4. AMA calls gun violence "a public health crisis" [news release]. Chicago, IL: American Medical Association; June 14, 2016. <https://www.ama-assn.org/ama-calls-gun-violence-public-health-crisis>. Accessed September 1, 2017.
5. Weinberger SE, Hoyt DB, Lawrence HC 3rd, et al. Firearm-related injury and death in the United States: a call to action from 8 health professional organizations and the American Bar Association. *Ann Intern Med*. 2015;162(7):513-516.
6. Physician groups representing 426,000 doctors: gun violence MUST STOP [news release]. Leawood, KS: American Academy of Family Physicians; June 14, 2016. <http://www.aafp.org/media-center/releases-statements/all/2016/violence-pub-health.html>. Accessed September 1, 2017.
7. Pinals DA, Appelbaum PS, Bonnie R, Fisher CE, Gold LH, Lee LW. American Psychiatric Association: position statement on firearm access, acts of violence and the relationship to mental illness and mental health services. *Behav Sci Law*. 2015;33(2-3):195-198.
8. Taichman DB, Laine C. Reducing firearm-related harms: time for us to study and speak out. *Ann Intern Med*. 2015;162(7):520-521.
9. American College of Surgeons. Statement on firearm injuries. <https://www.facs.org/about-ac/s/statements/12-firearm-injuries>. Published January 16, 2013. Accessed September 1, 2017.
10. American Trauma Society. Firearm violence. <http://www.amtrauma.org/?page=PosState132>. Published June 2013. Accessed September 1, 2017.
11. AMSA condemns gun violence, supports public health efforts moving forward [news release]. Sterling, VA: American Medical Student Association; June 15, 2017. <http://www.amsa.org/about/amsa-press-room/amsa-condemns-gun-violence-supports-public-health-efforts-moving-forward/>. Accessed September

- 1, 2017.
12. Wintemute GJ, Betz ME, Ranney ML. Yes, you can: physicians, patients, and firearms. *Ann Intern Med.* 2016;165(3):205-213.
 13. Betz ME, Wintemute GJ. Physician counseling on firearm safety: a new kind of cultural competence. *JAMA.* 2015;314(5):449-450.
 14. Barkin S, Duan N, Fink A, Brook RH, Gelberg L. The smoking gun: do clinicians follow guidelines on firearm safety counseling? *Arch Pediatr Adolesc Med.* 1998;152(8):749-756.
 15. Abraham A, Cheng TL, Wright JL, Addlestone I, Huang Z, Greenberg L. Assessing an educational intervention to improve physician violence screening skills. *Pediatrics.* 2001;107(5):e68.
<http://pediatrics.aappublications.org/content/107/5/e68.long>. Accessed November 22, 2017.
 16. Frank E, Carrera JS, Prystowsky J, Kellermann A. Firearm-related personal and clinical characteristics of US medical students. *South Med J.* 2006;99(3):216-225.
 17. Becher EC, Cassel CK, Nelson EA. Physician firearm ownership as a predictor of firearm injury prevention practice. *Am J Public Health.* 2000;90(10):1626-1628.
 18. Solomon BS, Duggan AK, Webster D, Serwint JR. Pediatric residents' attitudes and behaviors related to counseling adolescents and their parents about firearm safety. *Arch Pediatr Adolesc Med.* 2002;156(8):769-775.
 19. Betz ME, Barber CW, Miller M. Firearm restriction as suicide prevention: variation in belief and practice among providers in an urban emergency department. *Inj Prev.* 2010;16(4):278-281.
 20. Butkus R, Weissman A. Internists' attitudes toward prevention of firearm injury. *Ann Intern Med.* 2014;160(12):821-827.
 21. Price JH, Thompson AJ, Khubchandani J, Mrdjenovich AJ, Price JA. Firearm anticipatory guidance training in psychiatric residency programs. *Acad Psychiatry.* 2010;34(6):417-423.
 22. Khubchandani J, Price JH, Dake JA. Firearm injury prevention training in preventive medicine residency programs. *J Community Health.* 2009;34(4):295-300.
 23. Delnevo CD, Hausman AJ. Injury-prevention counseling among residents of internal medicine. *Am J Prev Med.* 2000;19(1):63-65.
 24. Giggie MA, Olvera RL, Joshi MN. Screening for risk factors associated with violence in pediatric patients presenting to a psychiatric emergency department. *J Psychiatr Pract.* 2007;13(4):246-252.
 25. Halpern-Felsher BL, Ozer EM, Millstein SG, et al. Preventive services in a health maintenance organization: how well do pediatricians screen and educate adolescent patients? *Arch Pediatr Adolesc Med.* 2000;154(2):173-179.
 26. Finch SA, Weiley V, Ip EH, Barkin S. Impact of pediatricians' perceived self-efficacy and confidence on violence prevention counseling: a national study. *Matern Child Health J.* 2008;12(1):75-82.
 27. Olson LM, Christoffel KK, O'Connor KG. Pediatricians' involvement in gun injury

- prevention. *Inj Prev*. 2007;13(2):99-104.
28. Puttagunta R, Coverdale TR, Coverdale J. What is taught on firearm safety in undergraduate, graduate, and continuing medical education? A review of educational programs. *Acad Psychiatry*. 2016;40(5):821-824.
 29. Pierson J, Viera AJ, Barnhouse KK, Tulsky JA, Richman BD, Goldstein AO. Physician attitudes and experience with permit applications for concealed weapons. *N Engl J Med*. 2014;370(25):2453-2454.
 30. Goldstein AO, Viera AJ, Pierson J, Barnhouse KK, Tulsky JA, Richman BD. Physician beliefs about physical and mental competency of patients applying for concealed weapon permits. *Behav Sci Law*. 2015;33(2-3):238-245.
 31. Parent B. Physicians asking patients about guns: promoting patient safety, respecting patient rights. *J Gen Intern Med*. 2016;31(10):1242-1245.
 32. Cassel CK, Nelson EA, Smith TW, Schwab CW, Barlow B, Gary NE. Internists' and surgeons' attitudes toward guns and firearm injury prevention. *Ann Intern Med*. 1998;128(3):224-230.
 33. Price JH, Kinnison A, Dake JA, Thompson AJ, Price JA. Psychiatrists' practices and perceptions regarding anticipatory guidance on firearms. *Am J Prev Med*. 2007;33(5):370-373.
 34. Price JH, Thompson A, Khubchandani J, Wiblishauser M, Dowling J, Teeple K. Perceived roles of emergency department physicians regarding anticipatory guidance on firearm safety. *J Emerg Med*. 2013;44(5):1007-1016.
 35. Kaplan MS, Adamek ME, Rhoades JA. Prevention of elderly suicide. Physicians' assessment of firearm availability. *Am J Prev Med*. 1998;15(1):60-64.
 36. Hethcoat GO 2nd. In the crosshairs: legislative restrictions on patient-physician speech about firearms. *DePaul J Health Care Law*. 2011;14(1):1-34.

Nicole D. Damari, MS, is a third-year medical student at the University of North Carolina School of Medicine in Chapel Hill. She holds an MS in pathobiology from Brown University. She is interested in population health, health policy, and the health of underserved and marginalized communities.

Karan S. Ahluwalia is a third-year medical student at the University of North Carolina School of Medicine in Chapel Hill. His interests include health policy, disparities, and prevention.

Anthony J. Viera, MD, MPH, is a professor in and the chair of the Department of Community and Family Medicine at Duke University School of Medicine in Durham, North Carolina, and an adjunct professor of public health leadership at the University of North Carolina Gillings School of Global Public Health. Among his interests is the role of policy in prevention of adverse health outcomes.

Adam O. Goldstein, MD, MPH, is a professor in the Department of Family Medicine at the University of North Carolina School of Medicine in Chapel Hill, where he also serves as the director of departmental advancement. Dr. Goldstein's research interests include policies and programs to prevent tobacco use, obesity, and firearm violence.

Acknowledgements

Research that informed this publication was supported by an institutional small grant from the University of North Carolina at Chapel Hill Department of Family Medicine.

Related in the *AMA Journal of Ethics*

[Law, Ethics, and Conversations between Physicians and Patients about Firearms in the Home](#), January 2018

[Physician "Gag Laws" and Gun Safety](#), April 2014

[Stop Posturing and Start Problem Solving: A Call for Research to Prevent Gun Violence](#), January 2018

[What Should Be the Scope of Physicians' Roles in Responding to Gun Violence?](#), January 2018

The viewpoints expressed in this article are those of the author(s) and do not necessarily reflect the views and policies of the AMA.

**Copyright 2018 American Medical Association. All rights reserved.
ISSN 2376-6980**